

MiR-488 inhibits proliferation and cisplatin sensibility in non-small-cell lung cancer (NSCLC) cells by activating the eIF3a-mediated NER signaling pathway


Chao Fang^{1,2}, Yi-Xin Chen^{1,2}, Na-Yiyuan Wu^{1,2}, Ji-Ye Yin^{1,2}, Xiang-Ping Li³, Hsuan-Shun Huang⁴, Wei Zhang^{1,2}, Hong-Hao Zhou^{1,2,5}, Zhao-Qian Liu^{1,2,5*}

Departments of Clinical Pharmacology¹ and Pharmacy³, Xiangya Hospital, Central South University, Changsha 410008, P. R. China; ²Institute of Clinical Pharmacology, Central South University, Hunan Key Laboratory of Pharmacogenetics, Changsha 410078, P. R. China;


⁴Department of Research, Cervical Cancer Prevention Center, Tzu Chi University, Hualien 970, Taiwan, Republic of China; ⁵Hunan Province Cooperation Innovation Center for Molecular Target New Drug Study, Hengyang 421001, P. R. China.

* Corresponding author : Professor Zhao-Qian Liu, Department of Clinical Pharmacology, Xiangya Hospital, Central South University, Changsha 410008; P. R. China; Institute of Clinical Pharmacology, Central South University; Hunan Key Laboratory of Pharmacogenetics, Changsha 410078; P. R. China.


Tel: +86 731 84805380, Fax: +86 731 82354476, E-mail: liuzhaoqian63@126.com.

(a)

H1299

(b)

SK-MES-1

(c)

A549

Figure S1. Transfection efficiency of miR-488 on eIF3a down-regulation.




Figure S2. The effect of miR-488 on the expression of P27 protein. Western blot was used to detect the P27 expression after the down-regulation of eIF3a with the miR-488 transfection.






Figure S3. Full-length blots